

Pollen morphology of some members of Piperaceae and its bearing on the systematics and phylogeny of the family

P. J. Mathew and P. M. Mathew*

Tropical Botanic Garden and Research Institute
Palode P.O., Thiruvananthapuram, 695 562, Kerala, India

*Perakathuseril, Muttada P.O.
Thiruvananthapuram, 695 025, Kerala, India

Abstract

Pollen morphology of 17 species of *Piper*, 8 cultivars of *Piper nigrum*, 13 species of *Peperomia* and the monotypic genus *Heckeria* of the family Piperaceae from South India was studied by both light and scanning electron microscopy. Two pollen types could be recognized : (1) *Piper* type (*Piper*, *Heckeria*) with isobilateral, monocolpate grains showing predominantly echinate exine surface pattern, and (2) *Peperomia* type (*Peperomia*) with radiosymmetric inaperturate grains showing predominantly areolate-spinulate exine. The pollen grains of all the three genera are very small-sized. Pollen features are shown to support closer affinity of the Piperaceae with Saururaceae. This also favours more evolved status of the *Piper* group than *Peperomia* within the family. The difference in the pollen features of *Peperomia* with the remainder of Piperaceae supports its segregation into a separate family Peperomiaceae.

INTRODUCTION

The Piperaceae are a large family comprising over 3000 species in 9 genera (Kupicha, 1993), of which *Piper* L. and *Peperomia* Ruiz & Pav. are the major ones, whose chief centres are Central and South America (Trelease & Yuncker, 1950). Their world distribution has been divided into 12 major centres (Datta & Dasgupta, 1977), one of which is the Indian subcontinent. In the Indian region, two independent centres of distribution have been recognized, namely the Trans-gangetic North-Eastern region and the South Deccan (Rahiman, 1987). Hooker (1886) described 56 species in 3 genera, viz., *Piper*, *Peperomia* and *Houttuynia* from the Indian subcontinent and Gamble (1925) recorded 13 species of *Piper*, 6 species of *Peperomia* and one in the monotypic genus *Heckeria* from South India. The family includes a number of economically and medicinally important species, of which *Piper nigrum* L. is the most important, which yields the 'Black pepper' of commerce, known as the 'King of Spices' a

P.J. Mathew and P.M. Mathew

highly valued condiment used as a culinary spice world over. Many species of *Peperomia* are favoured garden plants grown for their ornamental foliage.

Taxonomically the family is known to be a very puzzling group. Although palynological data have been recognized as a potential supplementary tool for tackling taxonomic and phylogenetic problems of related plant groups (Nair, 1974), palynological information on the Piperaceae is very scanty in general, and that too only on a handful of alien taxa (Erdtman, 1952; Smith, 1975). The South Indian group is almost unexplored. The present study concerns LM and SEM results of 17 species of *Piper*, 8 cultivars of *P. nigrum*, 13 species of *Peperomia* and one of *Heckeria* from South India, of which 4 species of *Piper* and 6 species of *Peperomia* are well established exotics grown as introduced plants in the region. The results of the study are discussed in relation to the systematics and phylogeny of the family.

MATERIALS AND METHODS

The polleniferous materials were collected from live plants from different localities of their distribution in South India. In the case of exotic species (*Piper arboreum*, *P. betle*, *P. colubrinum*, *P. magnificum*, *Peperomia clusiifolia*, *P. incana*, *P. obtusifolia*, *P. polybotrya*, *P. sandersii*, *P. scandens*) the materials used were from plants grown in the Tropical Botanic Garden and Research Institute, Palode, Thiruvananthapuram. Anthers from mature flowers were fixed in glacial acetic acid. Pollen preparations were made by acetolysis method (Erdtman, 1952) for LM study. For SEM study, the grains were stored in ethanol, coated with gold and scanned at the National Botanical Research Institute, Lucknow. Pollen terminology follows Erdtman (1967), and the pollen shape classes were determined after Walker and Doyle (1975). Pollen measurements were based on a sample size of 100 grains of each taxon. Voucher specimens of the taxa studied (Table 1) are deposited at the Herbarium of the Tropical Botanic Garden and Research Institute (TBGT), Palode, Thiruvananthapuram.

Table 1. List of taxa studied and their sources

Name of taxa	Voucher specimen No.	Locality/ Source	District & State
1	2	3	4
<i>Piper</i> L.			
<i>P. galeatum</i> C.DC.	19475	Munnar	Idukki, K
<i>P. trichostachyon</i> C.DC.	19426	Neymacadu	Idukki, K
<i>P. longum</i> L.	5848	Thenmala	Thiruvananthapuram, K
<i>P. hapnium</i> Buch. - Ham	19407	Palode	Thiruvananthapuram, K
<i>P. brachystachum</i> Wall.	8806	Kakki	Pathanamthitta, K
<i>P. hookeri</i> Miq.	8807	Kodaikanal	Anna, TN

Pollen morphology of some members of Piperaceae

1	2	3	4
<i>P. schmidtii</i> Hook.f.	8823	Avalanchi	Udagamandalam, TN
<i>P. barberi</i> Gamble	5699	Changili	Thiruvananthapuram, K
<i>P. nigrum</i> L. (cultivars)			
'Ampirian'	25333	Palode	Thiruvananthapuram, K
'Cherumany'	25315	Naiketty	Wayanad, K
'Kanjiramkoda'	25312	Kalpetta	Wayanad, K
'Karimunda'	12906	Peruvanthanam	Idukki, K
'Karimundy'	22621	Pulpally	Wayanad, K
'Kottanadan'	7737	Peringamala	Thiruvananthapuram, K
'Panickaruvally'	25314	Naiketty	Wayanad, K
'Peringamala'	25346	Peringamala	Thiruvananthapuram, K
<i>P. hymenophyllum</i> Miq.	7280	Pamba	Pathanamthitta, K
<i>P. argyrophyllum</i> Miq.	5023	Lower Kodayar	Tirunelveli, TN
<i>P. attenuatum</i> Miq.	2722	Athirapally	Trichur, K
<i>P. wightii</i> Miq.	8807	Agastymala	Thiruvananthapuram, K
<i>P. arboreum</i> Aubl.*	19454	TBGRI*	Thiruvananthapuram, K
<i>P. betle</i> L.*	19460	IIHR*	Bangalore, KA
<i>P. colubrinum</i> Link.*	12186	IISR*	Kozhikode, K
<i>P. magnificum</i> Trel.*	887	TBGRI	Thiruvananthapuram, K
<i>Heckeria</i> Kunth			
<i>H. subpeltata</i> Kunth [#]	13009	Palaruvi	Kollam, K
<i>Peperomia</i> Ruiz & Pav.			
<i>P. reflexa</i> A. Dietr.	12163	Athirumala	Thiruvananthapuram, K
<i>P. heyneana</i> Miq.	8831	Upper Kodayar	Tirunelveli, TN
<i>P. thomsoni</i> Hook.f.	19463	Ponmudi	Thiruvananthapuram, K
<i>P. portulacoides</i> A. Dietr.	12110	Peerumedu	Idukki, K
<i>P. wightiana</i> Miq.	19449	Periya	Wayanad, K
<i>P. dindigulensis</i> Miq.	12193	Adivaram	Kozhikode, K
<i>P. pellucida</i> H.B. & K.*	19470	TBGRI	Trivandrum, K
<i>P. clusiifolia</i> (Jacq.) Hook.*	12190	NGBS*	Wayanad, K
<i>P. incana</i> A. Dietr.	885	NGBS	Wayanadu, K

P.J. Mathew and P.M. Mathew

1	2	3	4
<i>P. obtusifolia</i> C.DC.*	19477	NGBS	Wayanad, K
<i>P. polybotrya</i> H.B. & K.*	19478	NGBS	Wayanad, K
<i>P. sandersii</i> C.DC. #	886	NGBS	Wayanad, K
<i>P. scandens</i> Ruiz. & Pav.*	19191	NGBS	Wayanad, K

* Exotic species

*TBGRI – Tropical Botanic Garden & Research Institute; *IIHR – Indian Institute of Horticultural Research; *IISR – Indian Institute of Spices Research; *NGBS – Narayana Gurukula Botanical Sanctuary; K – Kerala; KA – Karnataka; TN – Tamil Nadu.

RESULTS

The important pollen morphological features observed in species of *Piper*, *Peperomia* and *Heckeria* are summarized in Table 2. SEM pictures of the pollen of some species of *Piper* (Fig. 1: a-f; Fig. 2: g-l) and *Peperomia* (Fig. 3: m-r) are provided.

The pollen grains of all the species of *Piper*, and *Heckeria subpeltata* were consistently isobilateral and monocolpate with the aperture in the zonal position. The colpus furrow was narrow and long with tapering ends, almost extending to the poles in most species of *Piper*. However, in a few, the colpi were wider and shorter, and in certain cases (*P. schmidtii*) the ends of the colpus were broad and wide. All the species of *Peperomia* had radiosymmetric and inaperturate grains. The pollen grains of species of all the three genera were very small-sized, in general ranging from 6.2 x 6.0 µm – 9.0 x 6.66 µm; and in a few polyploid species of both *Piper* and *Peperomia* the grains were slightly larger; up to 11.3 x 9.8 µm in *Piper* (*P. brachystachyum*) and 13.2 x 10.4 µm in *Peperomia* (*P. portulacoides*). In one of the species of *Piper* (*P. galeatum*) the grains were extremely small sized (3.8 x 3.4 µm). The pollen shape-types noticed were spheroidal, prolate-spheroidal and subprolate in species of both *Piper* and *Peperomia*, and prolate-spheroidal in *Heckeria*. The predominant shape-type in *Piper* was prolate-spheroidal, while in *Peperomia* spheroidal. The exine surface patterns noticed in species of *Piper* were rugulate, verrucate, plated-spinulate or echinate, of which the echinate predominated. The surface in some of the species (see a, b - f in Fig.1 and i - l in Fig. 2) showed island / plate formations of various sizes and shapes. The surface in them is plated - spinulate, the plates being either at one level (layer) or at 2 or more levels (layers) superimposed over each other. Echinate condition is seen in 'b' and 'c' and in others spinulate. In *Peperomia* the most frequent type was the spinulate pattern, and in addition psilate, rugulate and echinate conditions were exhibited by one species each. In the species with spinulate pattern the spinules were seen to be located on differently sized and shaped areoles. The number of spinules per areole in different species showed wide variation (3-20); the highest number per areole was in *Peperomia dindigulensis* (14-20) followed by *P. polybotrya* (10-17),

Pollen morphology of some members of Piperaceae

and the lowest in *P. sandersii* (3-5). The projections are fundamentally tuberculate (m-o in Fig. 3) the tubercles being adpressed making a negative reticulum. The tubercles are free as in 'n' and adnate-tuberculate as in 'r' in Fig.3. The pollen shape in the species of *Heckeria* was prolate-spheroidal, and its exine surface was echinate.

Table 2. Important palynological features of species of the Piperaceae

Name of taxa	Aperture	Pollen shape	Pollen size (μ m)	Exine ornamentation
1	2	3	4	5
<i>Piper</i>				
<i>P. galeatum</i>	monocolpate	prolate – spheroidal	3.8 x 3.4	verrucate
<i>P. trichostachyon</i>	monocolpate	subprolate	8.4 x 6.4	spinulate
<i>P. longum</i>	monocolpate	prolate – spheroidal	8.1 x 6.9	spinulate
<i>P. hapnium</i>	monocolpate	prolate – spheroidal	7.5 x 6.5	spinulate
<i>P. brachystachyum</i>	monocolpate	subprolate	11.3 x 9.8	spinulate
<i>P. hookeri</i>	monocolpate	subprolate	7.1 x 6.9	granulate-rugulate
<i>P. schmidtii</i>	monocolpate	subprolate	7.3 x 5.9	spinulate
<i>P. barberi</i>	monocolpate	subprolate	8.3 x 6.3	echinate
<i>P. nigrum</i> (cultivars)				
'Ampirian'	monocolpate	subprolate	7.6 x 5.3	echinate
'Cherumany'	monocolpate	prolate – spheroidal	8.0 x 7.0	echinate
'Kanjiramkodan'	monocolpate	subprolate	6.8 x 5.4	areolate-spinulate
'Karimunda'	monocolpate	subprolate	7.6 x 5.4	echinate
'Karimundy'	monocolpate	spheroidal	6.2 x 6.0	echinate
'Kottanadan'	monocolpate	prolate – spheroidal	8.1 x 7.4	echinate
'Panickaruvally'	monocolpate	prolate – spheroidal	7.6 x 6.8	echinate
'Peringamala'	monocolpate	spheroidal	7.0 x 6.8	echinate
<i>P. hymenophyllum</i>	monocolpate	subprolate	7.8 x 6.5	echinate
<i>P. argyrophyllum</i>	monocolpate	prolate – spheroidal	7.2 x 6.8	echinate
<i>P. attenuatum</i>	monocolpate	prolate – spheroidal	10.0 x 9.0	echinate
<i>P. wightii</i>	monocolpate	subprolate	10.0 x 8.1	spinulate
<i>P. arboreum</i>	monocolpate	spheroidal	10.1 x 9.9	spinulate

P.J. Mathew and P.M. Mathew

1	2	3	4	5
<i>P. betle</i>	monocolpate	subprolate	7.3 x 5.4	echinate
<i>P. colubrinum</i>	monocolpate	prolate – spheroidal	9.0 x 6.6	spinulate
<i>P. magnificum</i>	monocolpate	subprolate	7.1 x 6.1	spinulate
Heckeria				
<i>H. subpeltata</i>	monocolpate	prolate-spheroidal	8.3 x 7.0	echinate
Peperomia				
<i>P. reflexa</i>	inaperturate	prolate-spheroidal	8.0 x 7.0	spinulate
<i>P. heyneana</i>	inaperturate	subprolate	7.6 x 6.0	areolate-spinulate
<i>P. thomsoni</i>	inaperturate	prolate-spheroidal	11.6 x 10.8	rugulate-perforate
<i>P. portulacoides</i>	inaperturate	subprolate	13.2 x 10.4	psilate
<i>P. wightiana</i>	inaperturate	prolate-spheroidal	7.2 x 6.8	areolate-spinulate
<i>P. dindigulensis</i>	inaperturate	prolate-spheroidal	8.3 x 7.8	areolate-spinulate
<i>P. pellucida</i>	inaperturate	prolate-spheroidal	8.9 x 8.0	areolate-spinulate
<i>P. clusifolia</i>	inaperturate	spheroidal	7.0 x 6.8	areolate-spinulate
<i>P. incana</i>	inaperturate	spheroidal	9.2 x 8.8	areolate-spinulate
<i>P. obtusifolia</i>	inaperturate	spheroidal	7.0 x 6.8	areolate-spinulate
<i>P. polybotrya</i>	inaperturate	spheroidal	6.8 x 6.7	areolate-spinulate
<i>P. sandersii</i>	inaperturate	prolate-spheroidal	7.0 x 6.4	areolate-spinulate
<i>P. scandens</i>	inaperturate	spheroidal	9.0 x 8.8	areolate-spinulate

Pollen morphology of some members of Piperaceae

DISCUSSION

Generally the cause of morphological evolution of pollen are assessed based on the position of the aperture, their number and form. Proximal, distal, zonal and global are considered to be the hierarchical order of evolutionary progress (Nair, 1974), of which the zonal and global are restricted to Angiosperms. Among the present taxa, members of the genus *Piper* and *Heckeria*, have their aperture in the zonal position. The number and distribution of the aperture forms are among the many trends in the evolution of pollen morphology (Walker & Doyle, 1975; Van Campo, 1976). An evolutionary sequence from simple to more complex apertural types has generally been contemplated (Chanda *et al.*, 1979), and increase in number from primitive to advanced appears to be the general trend. The pollen grains of monocots have one aperture and those of dicots usually three. The inaperturate (omniaperturate) condition occurs in the most primitive monocots (Helobiales) and primitive dicots (Polycarpae). In the present group, all the species of *Peperomia* are inaperturate and this is strikingly in contrast with the situation in the other two genera of the family, *Piper* and *Heckeria* in which the grains are consistently aperturate (monocolpate). As regards the size of the colpi, some degree of difference is apparent in *Piper*; some having long and narrow furrows with tapering ends and in others shorter and wider with round or broad colpus ends. Majority of the primitive families of the Ranalian complex, including the Saururaceae and Piperaceae are reported to show psilate exine sculpturing (cf. Walker, 1973) with scabrate, verrucate and echinate in addition. Among the present taxa of the family the most frequent exine surface pattern is echinate in *Piper* and also in *Heckeria*, while in *Peperomia* this is mostly spinulate.

The shape of pollen is usually unfixed in angiosperms, and hence this character has not been considered as a reliable parameter in pollen morphological analysis in relation to taxonomy and phylogeny. Judged by the P/E ratio, several shape classes are recognized in angiosperms (Walker & Doyle, 1975), of which the spheroidal is considered to be the most basic type, which changes in either direction. In the present group the shape categories noticed are spheroidal, prolate-spheroidal and subprolate, of which the prolate-spheroidal dominates in *Piper* and spheroidal in *Peperomia*. Pollen size is usually considered to be of less diagnostic value, but this character has been shown to be useful in cytopalynological studies (Bir & Sidhu, 1980; Nair & Ravikumar, 1984; Saraswathyamma *et al.*, 1995; Meenakumari *et al.*, 1996) in as much as the information on pollen size having provided an index to chromosome numerical variations. In the three Piperaceous genera studied here, the pollen grains are very small with size ranging from 6.2 x 6.0 – 11.3 x 9.8 μm in *Piper* and 6.8 x 6.7 - 13.2 x 10.4 μm in *Peperomia*. Although some of the polyploid taxa in both *Piper* and *Peperomia* such as the ($x=16$) *Piper brachystachyum* (Mathew *et al.* 1998 a) and ($x=4$) *Peperomia portulacoides* (Mathew *et al.*, 1998b) showed larger pollen size in relation to ploidy levels in both genera, there was no consistent positive correlation between pollen size and ploidy status.

In the light of the palynological data presented here on the South Indian members of the Piperaceae it may be possible to figure out two distinct pollen types viz., *Piper* type and *Peperomia* type in the family with the characteristics as shown below:

P.J. Mathew and P.M. Mathew

Pollen feature	Piper type (<i>Piper</i> , <i>Heckeria</i>)	Peperomia type (<i>Peperomia</i>)
a) Pollen symmetry	Isobilateral	Radiosymmetric
b) Pollen class	Monocolpate	Inaperturate
c) Exine surface	Predominantly echinate	Predominantly areolate-spinulate
d) Pollen shape	Predominantly prolate-spheroidal	Predominantly spheroidal

Pollen morphological data have been used in many studies related to cultivar taxonomy, and it has been demonstrated that statistical analysis of pollen variations has yielded viable information in the understanding of hybridity status of cultivars in crop species (Nair, 1961). In the South Indian State of Kerala there are a number of established and distinct varieties of *P. nigrum* under cultivation, of which pollen morphology of eight cultivars has been studied here. All the eight cultivars are similar in the basic aperture character (monocolpate), and their pollen size is also comparable. However, some difference was apparent in respect of the other features like pollen shape (spheroidal, prolate-spheroidal, subprolate). It is expected that a study of more cultivars of the species of the region, which is in progress, may yield meaningful data in relation to intraspecific taxonomy of this valuable crop species. Taxonomists are very much disagreed on many aspects of the systematics of the family, and there exists great confusion concerning the composition, interrelationships and affinities in the different classificatory treatments, both classical and modern. The family is considered to be a taxonomically difficult group (Burger, 1972) and is one of the most messes in plant taxonomy (Howard, 1973).

The members exhibit a number of anomalies, which puzzle any interpretation of plausible phylogenetic sequence. An important anatomical feature is that concerning the vascular bundles in two or more circles with the outer ones united or scattered as in monocots. This has prompted some of the early botanists to place it near the Araceae. The family also displays a suite of features, which are uncommon among dicots (Burger, 1977). But, unlike the monocots, the stem grows in thickness by cambial activity, which is a dicot feature (Metcalf & Chalk, 1950). Yuncker (1958) has held that embryonic and other features of dicot nature evident in the group more than offset the suspected monocot alliance. A significant attribute in relation to the phylogenetic position of the family is the simplicity of the floral structure. Whether the Piperaceous flower is to be reckoned as primitive or advanced is again a question on which there is no general concord. Concerning the affinities of the Piperaceae, there has been no agreement among taxonomists. Although affinity has been suggested with a number of primitive dicot families, most taxonomic treatments ascribe closer affinity with the Saururaceae (Bentham & Hooker, 1880; Johnson, 1902; Bessey, 1915; Skottsberg, 1946; Hutchinson, 1959; Thorne, 1976; Dahlgren, 1980; Takhtajan, 1980; Cronquist, 1981; Bedell & Reveal, 1982). Palynological semblance between the members of Saururaceae and *Piper* – the type genus of the Piperaceae (Walker, 1973) (very small, isobilateral, monocolpate grains with psilate, verrucate and echinate exine surface pattern) offer conceivable support to this affinity.

Pollen morphology of some members of Piperaceae




Fig. 1. Scanning electron micrographs of pollen grains of species of *Piper*: a. *P. arboreum* – exine spinulate; b. *P. betle* – exine echinate; c. *P. barberi* – exine echinate; d. *P. colubrinum*, exine spinulate; e. *P. brachystachyum* – exine spinulate; f. *P. hapnium* – exine spinulate.

P.J. Mathew and P.M. Mathew




Fig. 2. Scanning electron micrographs of pollen grains of species of *Piper*: g. *P. schmidtii* – exine spinulate; h. *P. longum* – exine spinulate; i. *P. nigrum* 'Kanjiramkoda' – exine areolate spinulate; j. *P. magnificum* – exine spinulate; k. *P. trichostachyon* – exine spinulate; l. *P. wightii* – exine spinulate.

Pollen morphology of some members of Piperaceae




Fig. 3. Scanning electron micrographs of pollen grains of species of *Peperomia*: m. *P. clusifolia* – exine areolate-spinulate with 4-6 spinules per areole; n. *P. incana* – exine areolate-spinulate with 7-12 spinules per areole; o. *P. polybotrya* – exine areolate-spinulate with 10-17 spinules per areole; p. *P. reflexa* – exine spinulate; q. *P. sandersii* – exine areolate-spinulate with 3-5 spinules per areole; r. *P. scandens* – exine areolate-spinulate with 5-7 spinules per areole.

P.J. Mathew and P.M. Mathew

The question of relative phylogenetic position of the two major genera, *Piper* and *Peperomia* within the family has been variously interpreted. The *Peperomia* group has been generally considered to be the more evolved and advanced for its herbaceous habit and more reduced floral structure. But, palynological features appear to subscribe more towards the view that the *Piper* group could be the more evolved in the family than *Peperomia* in as much as the former is endowed with aperturate (monocolpate) pollen grains with predominantly echinate exine surface as against the more primitive inaperturate grains with areolate-spinulate exine in *Peperomia*. The chromosomal features exhibited by the two genera also favour this possibility on the ground that the karyotypes of the species of *Piper* are consistently specialized with very small-sized chromosomes (Mathew *et al.*, 1998a) in contrast to the unspecialized karyotype of *Peperomia* in which the chromosomes are large-sized and meta- and submetacentric (Mathew *et al.*, 1998b).

Major similarities between *Piper* and *Peperomia* rest on the pattern of distribution of vascular bundles in the stem, spike-like inflorescence, bisexual flowers lacking perianth, one-celled ovary with one ovule. But *Peperomia* is distinct from *Piper* in respect of a spectrum of character-attributes like herbaceous habit, non-sheathing leaf base, lack of stipule, axillary spikes, more reduced flowers and also concerning the shape and appearance of subtending bracts (peltate-orbicular), position of stigma (eccentric), shape and size of fruit (ovoid, very small), etc. The wide exomorphic difference between the two generic groups has prompted some taxonomists to advocate segregation of the two, granting *Peperomia* the rank of a separate family, Peperomiaceae (Smith, 1972). The marked palynological distinction evident between *Piper* and *Peperomia* may be considered to offer convincing support to this proposition, which again is amply corroborated by the karyological distinction between the two genera.

Acknowledgement

The authors are thankful to Prof. G. Sreekandan Nair, Director, Tropical Botanic Garden and Research Institute, Palode, Thiruvananthapuram for providing facilities and encouragement.

Literature cited

- Bedell, H. G. & J. L. Reveal. 1982. Amended outlines and indices for six recently published systems of angiosperm classification. *Phytologia* 51: 65-156.
- Bentham, G. & J.D. Hooker. 1880. *Genera Plantarum*. L. Reeve & Co., London. Vol. I: 125-133.
- Bessey, C. E. 1915. The phylogenetic taxonomy of flowering plants. *Ann. Missouri Bot. Gard.* 2: 109-163.
- Bir, S. S. & M. Sidhu. 1980. Cytopalynological studies in weed flora of cultivated lands of Patiala District (Punjab). *J. Palynol.* 16: 85-105.

Pollen morphology of some members of Piperaceae

- Burger, W. C. 1972. The evolutionary trends in Central American species of *Piper*. *Brittonia* 24: 356-362.
- Burger, W. C. 1977. The Piperales and Monocots – Alternate hypothesis for the origin of monocotyledonous flowers. *Bot. Rev.* 43: 345-383.
- Chanda, S., K. Gosh & S. Nelson. 1979. On the polarity and tetrad arrangement in some mono- and diaperturate angiosperm pollen grains. *Grana* 18:21-31.
- Cronquist, A. 1981. *An Integrated System of Classification of Flowering plants*. Columbia Univ. Press, New York.
- Dahlgren, R.M. T. 1980. A revised system of classification of angiosperms. *Bot. J. Linn. Soc.* 80: 91-124.
- Datta P. C. & A. Dasgupta. 1977. Root anatomy and distribution of common *Piper* and *Peperomia* species. *Geobios* 4: 143 – 146.
- Erdtman, G. 1952. *Pollen morphology and plant taxonomy – Angiosperms*. Almqvist and Wiksell, Stockholm.
- Erdtman, G. 1967. *Handbook of Palynology*. Hafner Publishing Co., New York.
- Gamble, J. 1925. *Flora of the Presidency of Madras*. Adlard & Son Ltd., London.
- Hooker, J. D. 1886. *The Flora of British India*. L. Reeve & Co., London.
- Howard, R. A. 1973. Notes on the Piperaceae of Lesser Antilles. *J. Arnold Arb.* 54: 377-411.
- Hutchinson, J. 1959. *The Families of Flowering Plants*. Oxford University Press, London.
- Johnson, D.S. 1902. On the development of Piperaceae. *Bot. Gaz.* 34: 321–340.
- Kupicha, F. K. 1993. Piperaceae. In: V.H. Heywood (Ed.), *Flowering Plants of the World*. B.T. Batsford Ltd., London. pp. 38-46.
- Mathew, P. J., P. M. Mathew & P. Pushpangadan. 1998 a. Cytological studies in South Indian Piperaceae -I. *Piper*. *J. Cytol. Genet.* 33: 149 – 153.
- Mathew, P. J., P. M. Mathew & P. Pushpangadan. 1998 b. Cytological studies in South Indian Piperaceae- II. *Peperomia*. *J. Cytol. Genet.* 33: 155-158.
- Meenakumari, T., A. O. N. Panicker & C. K. Saraswathyamma. 1996. Pollen variability through induced polyploidy in *Pueraria phaseoloides* (Roxb.) Benth. *Palynol.* 32 : 85-93.
- Metcalf, C. R. & L. Chalk. 1950. *Anatomy of Dicotyledons*. The Clarendon Press, Oxford. pp. 1120-1127.
- Nair, P. K. K. 1961. Pollen grains in cultivated plants - II : *Bougainvillea*, *Hibiscus* and *Euphorbia*. *J. Ind. Bot. Soc.* 40: 365-381.

P.J. Mathew and P.M. Mathew

- Nair, P. K. K. 1974. Phylogeny of Angiosperms - II : A Palynological analysis. *Evol. Biol.* 7 : 127-137.
- Nair, P. K. K. & C. Ravikumar. 1984. Aspects and Prospects of Cytopalynology with special reference to pollen morphology. In: P.K.K. Nair (Ed.), *Advances in Pollen-Spore Research*. Today and Tomorrow's Printers and Publishers, New Delhi. pp.1-22.
- Rahiman, B. A. 1987. Geographical distribution of the genus *Piper* L. in India. *Curr. Sci.* 56 : 666
- Saraswathyamma, C. K., J. Licy & M. P. Sethuraj. 1995. Morphological and Palynological Investigations on diploids and tetraploids of para rubber tree *Hevea braziliensis* Muell.-Arg. *J. Palynol.* 31: 307-314.
- Skottsberg, G. 1946. *Peperomia berterona* Christoph.- An interesting case of disjunction. *Acta Hort. Gotob.* 16: 251 – 288.
- Smith, A. C. 1972. An appraisal of the orders and families of primitive extant families of Angiosperms. *J. Ind. Bot. Soc.* 50A: 215-229.
- Smith, C. A. 1975. The genus *Macropiper* (Piperaceae). *Bot. J. Linn. Soc.* 71: 1-38.
- Takhtajan, A. 1980. Outline of the classification of flowering plants. *Bot. Rev.* 46 : 225-359.
- Thorne, R. F. 1976. A phylogenetic classification of the angiospermae. *Evol. Biol.* 9: 35-106.
- Trelease, W. & T. G. Yuncker. 1950. *The Piperaceae of Northern South America*. University of Illinois Press, Urbana.
- Van Campo, M. 1976. Pattern of pollen morphological variation within taxa. In: Ferguson, I. K. & J. Muller (Eds.), *The evolutionary significance of the exine*. Academic Press, New York. pp. 163-183.
- Walker, J. W. 1973. Comparative pollen morphology and phylogeny of the ranalian complex. In: Charles B. Beck (Eds.), *Origin and early Evolution of Angiosperms*. Columbia University Press, New York. pp. 241-299.
- Walker, J. W. & J. A. Doyle. 1975. The bases of angiosperm phylogeny: Palynology. *Ann. Missouri Bot. Gard.* 62: 664-723.
- Yuncker, T. G. 1958. The Piperaceae – a family profile. *Brittonia* 10: 1-7.